




Listing of Claims

1. (Currently amended) A copolymer comprising at least one first monomeric unit and at least one second monomeric unit, wherein the at least one first monomeric unit has a Formulae I and I(a)




and the at least one second monomeric unit is selected from 5-membered-ring heteroaromatic groups having Formula IV



in each of Formulae I[, I(a)] and IV:

R is a substituent on a carbon atom which can be the same or different at each occurrence and is selected from hydrogen, alkyl, aryl, heteroalkyl, heteroaryl, F, -

CN, $-OR^1$, $-CO_2R^1$, $-C_{\psi}H_{\theta}F_{\lambda}$, $-OC_{\psi}H_{\theta}F_{\lambda}$, $-SR^1$, $-N(R^1)_2$, $-P(R^1)_2$, $-SOR^1$, $-SO_2R^1$, $-NO_2$, and beta-dicarbonyls having Formula XII



or adjacent R groups together can form a 5- or 6-membered cycloalkyl, aryl, or heteroaryl ring,

such that:

R^1 is a substituent on a heteroatom which can be the same or different at each occurrence and is selected from alkyl, aryl, heteroalkyl and heteroaryl; and ψ is an integer between 1 and 20, and θ and λ are integers satisfying Equation A1 below:

$$\theta + \lambda = 2\psi + 1; \quad (\text{Equation A1});$$

in Formula IV:

E can be the same or different at each occurrence and is a single bond or a linking group selected from arylene and heteroarylene;

in Formula IV:

A is independently at each occurrence C or N and γ is 0 or an integer selected from 1 or 2, such that when both A are N, then γ is 0; or when one of A is N and one of A is C, then γ is 1; or when both A are C, then γ is 2;

Q is O, S, SO_2 , or NR^1 where:

R^1 is a substituent on a heteroatom which can be the same or different at each occurrence and is selected from alkyl, aryl, heteroalkyl and heteroaryl;

in Formula XII:

R^2 is selected from hydrogen, alkyl, aryl, heteroalkyl and heteroaryl;

δ is 0 or an integer from 1 to 12, and when R in formula IV is hydrogen, alkyl, F, $-CN$, $-OR^1$, or CO_2R^1 the copolymer further comprises end-capping groups that are aromatic.

Application No.: 10/809,142
 Docket No.: PE0667 US DIV1

Page 4

2. (Original) The copolymer of Claim 1, wherein R groups in one or more of the at least one first monomeric unit are independently selected from alkyl groups having 1 to 30 carbon atoms; heteroalkyl groups having 1-30 carbon atoms and one or more heteroatoms of S, N, or O; aryl groups having from 6 to 20 carbon atoms, and heteroaryl groups having from 2 to 20 carbon atoms and one or more heteroatoms of S, N, or O.

3. (Original) The copolymer of Claim 1 that excludes any vinylene monomeric units.

4. (Previously presented) The copolymer of Claim 1 wherein each R group in each of Formula I, Formula I(a), and Formula IV is selected from:

hydrogen;

alkyl;

aryl;

heteroalkyl;

heteroaryl;


F;

-CN;

-P(R¹)₂ and -SOR¹, where R¹ is a substituent on a heteroatom which can be the same or different at each occurrence and is selected from alkyl, aryl, heteroalkyl and heteroaryl;

-NO₂;

a beta-dicarbonyl having Formula XII



-C_ψH_θF_λ;

-OC_ψH_θF_λ;

-OR¹, -CO₂R¹, -SR¹, -N(R¹)₂, and -SO₂R¹ where R¹ is a straight chain or branched alkyl of more than 20 carbons or a straight chain or branched heteroalkyl.


Application No.: 10/809,142
Docket No.: PE0667 US DIV1

Page 5

5. (Original) The copolymer of Claim 1 wherein the at least one of the R groups in one or more of the at least one first monomeric unit is independently selected from linear and branched n-butyl groups; linear and branched iso-butyl groups; linear and branched pentyl groups; hexyl groups, and octyl groups with and without olefinic unsaturation; phenyl groups, thiophene groups, carbazole groups, alkoxy groups, phenoxy groups and cyano groups.

6. (Original) The copolymer of Claim 1 wherein at least one of the R groups in one or more of the at least one first monomeric unit are independently selected from H, C₆-C₁₂ alkoxy, phenoxy, C₆-C₁₂ alkyl, phenyl and cyano.

7. (Currently Amended) The copolymer of Claim 1 wherein one or more of the at least one second monomeric unit is selected from Formulae I, I(a), and IV(a) through IV(h):



Application No.: 10/809,142
Docket No.: PE0667 US DIV1

Page 6

where:

in Formula IV(a):

R is as described above for each of Formulae I, I(a) and IV;

in Formula IV(h):

R¹ is a substituent on a heteroatom which can be the same or different at each occurrence and is selected from alkyl, aryl, heteroalkyl and heteroaryl.

8. (Cancelled).

9. (Previously Presented) The copolymer of Claim 1, wherein one or more of the at least one second monomeric unit has Formula IV wherein R is selected from:

partially or fully fluorinated alkyl groups having from 1 to 12 carbon atoms;


alkoxy groups having from 1 to 12 carbon atoms;

esters having from 3 to 15 carbon atoms;

-SR¹, -N(R¹)₂, -P(R¹)₂, -SOR¹, -SO₂R¹, where R¹ is an alkyl group having from 1 to 12 carbon atoms;

-NO₂; and

beta-dicarbonyls having Formula XII where:



in Formula XII:

R² is an alkyl group having from 1 to 12 carbon atoms and δ is 0, 1, or 2.

10. (Cancelled).

11. (Original) The copolymer of Claim 1, wherein one or more of the at least one second monomeric unit has Formula IV wherein:

R groups are selected from H, C₆-C₁₂ alkyl groups, C₆-C₂₀ aryl groups, and C₂-C₂₀ heteroaryl groups; and

E linking groups include pyrrolediyl (-C₄H₃N-) and thiophenediyl (-C₄H₃S-).

Application No.: 10/809,142
Docket No.: PE0667 US DIV1

Page 7

12-13. (Cancelled).

14. (Original) An electronic device comprising at least one electroactive layer comprising the copolymer of Claim 1.

15. (Original) The device of Claim 14, wherein the device comprises a hole injection/transport layer comprising the copolymer of Claim 1.

16. (Original) The device of Claim 14, wherein the device comprises an electron injection/transport layer comprising the copolymer of Claim 1.

17. (Original) The device of Claim 14, wherein the electroactive layer comprises a light-emitting material comprising the copolymer of Claim 1.


18. (Cancelled).

19. (Original) The device of Claim 14, wherein the device is selected from a light-emitting device, a photodetector, and a photovoltaic device.

20. (Original) The device of Claim 14, wherein the device is an electroluminescent display.


21. (Currently Amended) A light-emitting device comprising at least one light-emitting layer comprising the following copolymer;

at least one first monomeric unit and at least one second monomeric unit, wherein the at least one first monomeric unit has a Formulae I and I(a)




Application No.: 10/809,142
 Docket No.: PE0667 US DIV1

Page 8




and the at least one second monomeric unit is selected from 5-membered-ring heteroaromatic groups having Formula IV



in each of Formulae I[[, I(a)]] and IV:

R is a substituent on a carbon atom which can be the same or different at each occurrence and is selected from hydrogen, alkyl, aryl, heteroalkyl, heteroaryl, F, -CN, -OR¹, -CO₂R¹, -C_ψH_θF_λ, -OC_ψH_θF_λ, -SR¹, -N(R¹)₂, -P(R¹)₂, -SOR¹, -SO₂R¹, -NO₂, and beta-dicarbonyls having Formula XII



or adjacent R groups together can form a 5- or 6-membered cycloalkyl, aryl, or heteroaryl ring,

such that:

R¹ is a substituent on a heteroatom which can be the same or different at each occurrence and is selected from alkyl, aryl, heteroalkyl and heteroaryl; and

Application No.: 10/809,142
Docket No.: PE0667 US DIV1

Page 9

ψ is an integer between 1 and 20, and θ and λ are integers satisfying Equation A1 below:

$$\theta + \lambda = 2\psi + 1; \quad (\text{Equation A1});$$

in Formula IV:

E can be the same or different at each occurrence and is a single bond or a linking group selected from arylene and heteroarylene;

in Formula IV:

A is independently at each occurrence C or N and γ is 0 or an integer selected from 1 or 2, such that when both A are N, then γ is 0; or when one of A is N and one of A is C, then γ is 1; or when both A are C, then γ is 2;

Q is O, S, SO₂, or NR¹ where:

R¹ is a substituent on a heteroatom which can be the same or different at each occurrence and is selected from alkyl, aryl, heteroalkyl and heteroaryl [(.)] _ _

in Formula XII:

R² is selected from hydrogen, alkyl, aryl, heteroalkyl and heteroaryl;

δ is 0 or an integer from 1 to 12.